

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07D 215/56, 215/22, 215/18, A61K 31/47</b>		(11) International Publication Number: <b>WO 00/03991</b>
A1		(43) International Publication Date: 27 January 2000 (27.01.00)
(21) International Application Number: PCT/SE99/01270		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 14 July 1999 (14.07.99)		
(30) Priority Data: 9802549-7 15 July 1998 (15.07.98) SE		
(71) Applicant: ACTIVE BIOTECH AB [SE/SE]; P.O. Box 724, S-220 07 Lund (SE).		
(72) Inventors: BJÖRK, Anders; Svalvågen 9, S-237 36 Bjärred (SE). JÖNSSON, Stig; Sofiaparken 2, S-222 41 Lund (SE). FEX, Tomas; Harpasset 1, S-226 52 Lund (SE). HEDLUND, Gunnar; Gullregnsvägen 131 E, S-224 56 Lund (SE).		
(74) Agents: BERGVALL EFTRING, Stina, Lena et al.; Dr. Ludwig Brann Patentbyrå AB, P.O. Box 17192, S-104 62 Stockholm (SE).		Published With international search report.

(54) Title: QUINOLINE DERIVATIVES



(57) Abstract

The invention is related to compounds of general formula (I), wherein R is methyl, ethyl, n-propyl, iso-propyl, n-butyl or allyl; R' is methyl, methoxy, fluoro, chloro, bromo, trifluoromethyl, or OCH<sub>2</sub>F<sub>y</sub>, wherein x = 0 - 2, y = 1 - 3 with the proviso that x + y = 3; R'' is hydrogen, fluoro or chloro; with the proviso that R'' is fluoro or chloro only when R' is fluoro; R<sub>4</sub> is hydrogen or pharmaceutically acceptable inorganic or organic cations; R<sub>5</sub> is ethyl, n-propyl, iso-propyl, methoxy, ethoxy, chloro, bromo, trifluoromethyl, OCH<sub>2</sub>F<sub>y</sub>, or OCH<sub>2</sub>CH<sub>2</sub>F<sub>y</sub>, wherein x = 0 - 2, y = 1 - 3 with the proviso that x + y = 3; R<sub>6</sub> is hydrogen; or R<sub>5</sub> and R<sub>6</sub> taken together are methylenedioxy; and any tautomer thereof. The invention also relates to pharmaceutical compositions containing a compound of general formula (I) together with a pharmaceutically acceptable carrier. Included are also processes for the preparation of the compounds of formula (I), as well as methods for the treatment of mammals suffering from diseases resulting from autoimmunity and pathological inflammation by administering of a compound having formula (I) to said mammal.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## QUINOLINE DERIVATIVES

### FIELD OF THE INVENTION

The present invention relates to novel substituted quinoline-3-carboxamide derivatives, to methods for their preparation, to compositions containing them, and to methods and use for clinical treatment of diseases resulting from autoimmunity, such as multiple sclerosis, insulin-dependent diabetes mellitus, systemic lupus erythematosus, rheumatoid arthritis, inflammatory bowel disease and psoriasis and, furthermore, diseases where pathologic inflammation plays a major role, such as asthma, atherosclerosis, stroke and Alzheimer's disease. More particularly, the present invention relates to novel quinoline derivatives suitable for the treatment of, for example, multiple sclerosis and its manifestations.

### BACKGROUND OF THE INVENTION

Autoimmune diseases, e.g., multiple sclerosis (MS), insulin-dependent diabetes mellitus (IDDM), systemic lupus erythematosus (SLE), rheumatoid arthritis (RA), inflammatory bowel disease (IBD) and psoriasis represent assaults by the body's immune system which may be systemic in nature, or else directed at individual organs in the body. They appear to be diseases in which the immune system makes mistakes and, instead of mediating protective functions, becomes the aggressor (1).


MS is the most common acquired neurologic disease of young adults in western Europe and North America. It accounts for more disability and financial loss, both in lost income and in medical care, than any other neurologic disease of this age group. There are approximately 250,000 cases of MS in the United States.

Although the cause of MS is unknown, advances in brain imaging, immunology, and molecular biology have increased researchers' understanding of this disease. Several therapies are currently being used to treat MS, but no single treatment has demonstrated dramatic

treatment efficacy. Current treatment of MS falls into three categories: treatment of acute exacerbations, modulation of progressive disease, and therapy for specific symptoms.

MS affects the central nervous system and involves a demyelination process, i.e., the myelin sheaths are lost whereas the axons are preserved. Myelin provides the isolating material that enables rapid nerve impulse conduction. Evidently, in demyelination, this property is lost. Although the pathogenic mechanisms responsible for MS are not understood, several lines of evidence indicate that demyelination has an immunopathologic basis. The pathologic lesions, the plaques, are characterised by infiltration of immunologically active cells such as macrophages and activated T cells (2).

In US Pat. No. 4,547,511 and in US Pat. No. 4,738,971 and in EP 59,698 some derivatives of N-aryl-1,2-dihydro-4-substituted-1-alkyl-2-oxo-quinoline-3-carboxamide are claimed as enhancers of cell-mediated immunity. The compound



Roquinimex

known as roquinimex (Merck Index 12th Ed., No. 8418; Linomide<sup>®</sup>, LS2616, N-phenyl-N-methyl-1,2-dihydro-4-hydroxy-1-methyl-2-oxo-quinoline-3-carboxamide) belongs to this series of compounds. Roquinimex has been reported to have multiple immunomodulatory activities not accompanied with general immunosuppression (3-12).

Furthermore, in US Pat. No. 5,580,882 quinoline-3-carboxamide derivatives are claimed to be useful in the treatment of conditions associated with MS. The particular preferred compound is roquinimex. In US Pat. No. 5,594,005 quinoline-3-carboxamide derivatives are claimed to be useful in the treatment of type I diabetes. The particular preferred compound is roquinimex.

In WO 95/24195 quinoline-3-carboxamide derivatives are claimed to be useful in the treatment of inflammatory bowel disease. Particularly preferred compounds are roquinimex or a salt thereof. In WO95/24196 quinoline-3-carboxamide derivatives are claimed to be useful in the treatment of psoriasis. Particularly preferred compounds are roquinimex or a salt thereof.

In clinical trials comparing roquinimex to placebo, roquinimex was reported to hold promise in the treatment of conditions associated with MS (13, 14). There are, however, some serious drawbacks connected to roquinimex. For example, it has been found to be teratogenic in the rat, and to induce dose-limiting side effects in man, e.g., a flu-like syndrome, which prevents from using the full clinical potential of the compound.

Further, in WO 92/18483 quinoline derivatives substituted in the 6-position with a  $R_A S(O)_n$ -group ( $R_A$  = lower alkyl or aryl;  $n = 0 - 2$ ) are claimed, which possess an immunomodulating, anti-inflammatory and anti-cancer effect.

The substitution, i.e., type and pattern, of the above, specifically mentioned, compounds in the prior art places them outside the scope of the present invention.


#### DESCRIPTION OF THE INVENTION

A primary objective of the present invention is to provide structurally novel quinoline compounds which by virtue of their pharmacological profile, with high potency in experimental models and low level of side-effects, are considered to be of value in the treatment of disease resulting from autoimmunity and pathologic inflammation. Examples of such diseases are multiple sclerosis, insulin-dependent diabetes mellitus, systemic lupus erythematosus, rheumatoid arthritis, inflammatory bowel disease and psoriasis and other diseases where inflammation plays a major role, such as asthma, atherosclerosis, stroke and Alzheimer's disease. More particularly, the present invention relates to novel quinoline derivatives suitable for the treatment of, for example, multiple sclerosis and its manifestations.

The term "treatment" as used herein includes prophylaxis as well as relieving the symptoms of

disease.

It has now surprisingly been found that the novel compounds of general formula (I)



wherein

R is selected from methyl, ethyl, n-propyl, iso-propyl, n-butyl and allyl;

R' is selected from methyl, methoxy, fluoro, chloro, bromo, trifluoromethyl, and  $\text{OCH}_x\text{F}_y$ ,

wherein  $x = 0 - 2$ ,

$y = 1 - 3$  with the proviso that

$x + y = 3$ ;

R'' is selected from hydrogen, fluoro and chloro, with the proviso that R'' is selected from fluoro and chloro only when R' is selected from fluoro and chloro;

R<sub>4</sub> is selected from hydrogen and pharmaceutically acceptable inorganic cations, such as sodium, potassium and calcium, and organic cations such as monoethanolamine, diethanolamine, dimethylaminoethanol, morpholine and the like;

R<sub>5</sub> is selected from ethyl, n-propyl, iso-propyl, methoxy, ethoxy, chloro, bromo, trifluoromethyl, and  $\text{OCH}_x\text{F}_y$ , and  $\text{OCH}_2\text{CH}_x\text{F}_y$ ,

wherein  $x = 0 - 2$ ,

$y = 1 - 3$  with the proviso that

$x + y = 3$ ;

R<sub>6</sub> is hydrogen; or

R<sub>5</sub> and R<sub>6</sub> taken together are methylenedioxy;

are unexpectedly effective and specific in the treatment of individuals suffering from autoimmune and inflammatory diseases.

The compounds of general formula (I) may exist in different tautomeric forms and all such forms where such forms exist are included herein.

In a preferred embodiment of the invention  $R_4$  is selected from hydrogen or sodium,

$R_5$  is selected from ethyl, methoxy, chloro and trifluoromethyl,

$R_5$  and  $R_6$  taken together are methylenedioxy,

$R$  is selected from methyl and ethyl,

$R'$  is selected from methoxy, fluoro, chloro and trifluoromethyl when  $R''$  is hydrogen and

$R''$  is selected from *meta*- and *para*-fluoro provided that  $R'$  is *ortho*-fluoro.

Among the most preferred compounds of general formula (I) according to the present invention are:

N-ethyl-N-(3-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

N-ethyl-N-(4-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

N-ethyl-N-(3-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-ethyl-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(2-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(4-chloro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,  
N-methyl-N-(4-trifluoromethyl-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,  
N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5,6-methylenedioxy-1-methyl-2-oxo-quinoline-3-carboxamide,  
N-methyl-N-(4-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-trifluoromethyl-1-methyl-2-oxo-quinoline-3-carboxamide.

Several spontaneously occurring autoimmune diseases in man have experimental models that are spontaneously occurring in certain strains of laboratory animals or can be induced in laboratory animals by immunisation with specific antigen(s) from the target organ.

Experimental autoimmune encephalomyelitis (EAE) as a model for autoimmune inflammatory diseases of the central nervous system (CNS) has been the most widely used model for the human disease multiple sclerosis.

Autoimmunity to type II collagen can experimentally be induced in certain strains of mice or rats and may lead to the development of polyarthritis. The collagen induced arthritis has several features in common with the human disorder rheumatoid arthritis.

The hallmark of asthma in humans is an increased reactivity of the airways to a range of chemical and physical stimuli. It is now widely accepted that products released from inflammatory cells, e.g., activated eosinophils, compromise epithelial integrity and promote bronchial hyperresponsiveness. The murine model of ovalbumin (OA)-induced lung inflammation is dominated by the temporally regulated influx of lymphocytes and eosinophils into the bronchial lumen.

Roquinimex has been found to induce the Beagle Pain Syndrome (BPS) (15, 16) in different breeds of beagle dogs. The disease is reflected by clinical and laboratory manifestations justifying BPS as a model for the flu-like syndrome induced by roquinimex in man.




The compounds of general formula (I) were assayed for inhibition of acute experimental autoimmune encephalomyelitis (aEAE) in mice. Roquinimex was used as treatment control and showed a more than 50% inhibition at  $\geq 5$  mg/kg. Surprising and unexpected results were obtained when introducing proper substitution in the 5-position of the quinoline ring. In comparison with roquinimex, the potency of the 5-chloro substituted compound was increased a 100-fold. Substitution in the 6-, 7-, and 8-position resulted in less active compounds. The effect of the 5-substitution could largely be understood on physicochemical grounds. In general, the EAE activity as seen by the EAE inhibition was in the following descending order according to the position of the substitution:  $5 > 6 \gg 7 = 8$ . The comparison of the effects of 5- and 6-substitution showed that there is a statistically significant difference on every normal level ( $p < 0.001$ ) between the two, the effect of the 5-substitution being superior. Furthermore, proper aromatic substitution in the quinoline moiety and the 3-carboxamide moiety of the compounds of general formula (I) significantly reduced or even abolished the side-effects, i.e., the teratogenic effect and the BPS, of roquinimex. Thus, physicochemical properties of the 5-substituent in the quinoline moiety and the *ortho*-, *meta*- and/or, in particular, the *para*-substituent in the 3-carboxamide moiety are of major importance for an improved risk/benefit ratio in comparison with roquinimex. Replacement of the methyl group on the carboxamide nitrogen with a higher alkyl group reduced the side effects even further. Hence, the compounds of formula (I) have surprisingly been found to be both chemically and pharmacologically different from those drugs hitherto suggested for the treatment of MS and its manifestations.


All embodiments of the invention as disclosed in the claims are herewith included in the specification.

The following examples are intended to illustrate the invention without restricting the scope thereof.

The compounds of general formula (I) may be prepared by methods known in the literature and the following methods:

Method A:

The compounds of general formula (I) may be prepared by known methods and, for example, as shown above, by reaction of an ester derivative of the quinoline carboxylic acid with an aniline in a suitable solvent such as toluene, xylene and the like. General methods for preparation of the quinoline carboxylic acid ester derivatives of formula (II) are described below. N-alkylated anilines of formula (III) are commercially available or known from literature, e.g., in Johnstone et al, J. Chem. Soc. 1969, 2223-2224. Compounds falling within the scope of formula (III) may be prepared by methods, which are generally analogous to those of said literature.

Method B:

The compounds of formula (I) may also be prepared by reaction of a quinoline carboxylic acid of formula (IV) with an aniline of formula (III). Various coupling reagents known in the art may be used, e.g., carbodiimides known from literature in US Pat. No. 4,547,511. One suitable coupling method utilises thionyl chloride in the presence of triethylamine and a suitable solvent such as dichloromethane. This method may be used in instances when direct coupling

between ester and aniline does not work, e.g., when the aniline contains electron withdrawing substituents. The quinoline carboxylic acids of formula (IV) may be obtained from the corresponding esters of formula (II) by acidic hydrolysis as described below.

*Example 1.*

1,2-Dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxylic acid ethyl ester.

To a solution of 2,6-difluorobenzonitril (42 g, 0.30 mol) in 150 ml of anhydrous methanol sodium methoxide (17.9 g, 0.33 mol) was slowly added at 30°C. After being heated under reflux for 1 hour, aqueous 40 % methylamine (133 ml, 1.2 mol) was added and the resulting solution refluxed for 4 days. On cooling, a white solid, 2-methoxy-6-(methylamino)benzonitrile, precipitated which was collected by filtration. The precipitate was dissolved in an aqueous solution of ethylene glycol (500 ml) and potassium hydroxide (14 g). The solution was refluxed at 150°C over night, cooled to room temperature and the pH adjusted to 4 with conc. hydrochloric acid. The anthranilic acid formed was collected by filtration, washed with water (50 ml) and dried under vacuum. The 6-methoxy-N-methyl-anthranilic acid (32 g, 0.18 mol), and sodium bicarbonate (38 g, 0.45 mol) were suspended in 500 ml of 1,4-dioxane. Phosgene (25 ml, 0.45 mol) was slowly added under cooling in an ice bath. The mixture was warmed at 40°C for 1 hour, cooled to 15°C, and then 150 ml of water was added. The isatoic anhydride formed was collected by filtration. After being carefully dried, the 5-methoxy-N-methyl-isatoic anhydride (20.7 g, 0.10 mol) was added to a solution of sodium diethylmalonate (31 g, 0.17 mol) in 250 ml of anhydrous N,N-dimethylformamide at room temperature. The solution was heated at 100°C for 3 hours, cooled to room temperature, 250 ml of water was added and the pH adjusted to 4 with conc. hydrochloric acid. The precipitate was collected by filtration and dried in vacuum to give the title compound as pure white crystals (22 g), yield 48 %.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.43 (t, 3H), 3.62 (s, 3H), 3.96 (s, 3H), 4.45 (q, 2H), 6.70 (d, 1H), 6.92 (d, 1H), 7.55 (t, 1H), 13.5 (s, 1H).

*Example 2.*

1,2-Dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxylic acid ethyl ester.

Phosgene (51 g, 0.52 mol) dissolved in 150 ml of dioxane was added in portions to a

mechanically stirred slurry of 2-amino-6-chloro-benzoic acid (30 g, 0.175 mol) and sodium bicarbonate (44 g, 0.52 mol) in 300 ml of dioxane. Violent reaction with gas evolution occurred and the reaction mixture was cooled to keep the temperature below 50°C. Then stirring was continued at 50°C for 1 hour. The reaction mixture was cooled to 15°C, the resulting precipitate was collected, washed with water and dried to give the isatoic anhydride. The anhydride (5 g, 0.025 mol) was dissolved in 50 ml of N,N-dimethylacetamide and cooled to 0°C. Sodium hydride (75%) (0.94 g, 0.028 mol) and then methyl iodide (1.89 ml, 0.030 mol) was added at a rate to keep the temperature below 5°C. The reaction mixture was stirred at room temperature for 5 hours. The remaining methyl iodide was removed under vacuum. Sodium hydride (0.94 g, 0.028 mol) was added together with diethyl malonate (4.5 g, 0.028 mol). The mixture was heated at 85°C for 5 hours. After cooling to room temperature, 50 ml of methanol and 50 ml of 1 M hydrochloric acid and subsequently 250 ml of water were added. An emulsion was formed which crystallised on standing in a refrigerator for 72 hours. The crystalline mass was collected by filtration, washed with water, water/methanol (1:1) and heptane and dried to afford the title compound (6.3 g), yield 74 %.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.46 (3H, t), 3.63 (3H, s), 4.49 (2H, q), 7.23 (1H, d), 7.27 (1H, d), 7.49 (1H, t), 15.0 (1H, s).

### *Example 3.*

#### 1,2-Dihydro-4-hydroxy-5-trifluoromethyl-1-methyl-2-oxo-quinoline-3-carboxylic acid ethyl ester.

2-Fluoro-6-(trifluoromethyl)benzonitrile (10 g, 0.053 mol) was warmed at 40°C in 200 ml of anhydrous methylamine in an autoclave for 2 days. The excess methylamine was allowed to evaporate and the resulting grey solid was dissolved in 200 ml of methylene chloride together with 4-aminopyridine (0.1 g, 0.001 mol) and triethylamine (3.3 ml, 0.026 mol). To the chilled solution was slowly added ethyl malonyl chloride (8.8 g, 0.060 mol). The solution was stirred for 4 hours and then worked up to give a yellowish syrup. The syrup was dissolved in 100 ml of anhydrous ethanol, and sodium methoxide (5.4 g, 0.10 mol) was added. After 1 hour, the solvent was removed and the residue worked up with methylene chloride and water. The quinoline derivative formed was carefully dried and suspended in 250 ml of chilled anhydrous tetrahydrofuran. Sodium hydride (4 g, 0.125 mol) was slowly added and then methyl iodide

(10 ml, 0.15 mol). The mixture was heated under reflux for 6 hours, quenched with water and worked up with diethyl ether. The solvents were removed and the residue (17.3 g) was dissolved in a mixture of ethanol (50 ml) and conc. hydrochloric acid (10 ml). The solution was warmed at 45°C overnight, cooled and the precipitate was collected to give 8 g of the title compound, yield 48 %.

<sup>1</sup>H NMR  $\delta$  (CDCl<sub>3</sub>)  $\delta$  1.46 (3H, t), 3.68 (3H, s), 4.50 (2H, q), 7.58 (1H, m), 7.71 (2H, m), 15.0 (1H, s).

In essentially the same manner the following compound was obtained from the corresponding starting materials:

1,2-Dihydro-4-hydroxy-5-trifluoromethoxy-1-methyl-2-oxo-quinoline-3-carboxylic acid ethyl ester.

*Example 4.*

1,2-dihydro-4-hydroxy-1-methyl-2-oxo-5,6-methylenedioxy-quinoline-3-carboxylic acid ethyl ester.

Di-tert-butyl dicarbonate (36 g, 0.17 mol) was added portionwise to a solution of 3,4-(methylenedioxy)-aniline (20.6 g, 0.15 mol) in anhydrous tetrahydrofuran (150 ml). The solution was reflux heated for 2 hours, then concentrated under vacuum to give a black solid residue. The residue was dissolved in anhydrous tetrahydrofuran (600 ml) and cooled to -40°C. A hexane solution of 1.3 M *sec*-butyllithium (265 ml, 0.35 mol) was added dropwise. After stirring the solution for 0.5 hour at -40°C dry ice (ca 40 g) pellets were added. The mixture was allowed to warm to 0°C and water (ca 700 ml) was added. The aqueous solution was acidified with hydrochloric acid to pH 3 and extracted with ether. The extracts were dried and concentrated to give the N-tBoc protected 5,6-methylenedioxy-anthranilic acid as a solid residue (45 g). This acid was added to an ice-cooled suspension of sodium hydride (80 % in oil, 9.0 g, 0.30 mol) in N,N-dimethylformamide (200 ml). The mixture was stirred for 0.5 hour and methyl iodide (22 ml, 0.35 mol) was added. The mixture was stirred at room temperature overnight, was quenched with water (600 ml) and extracted three times with ether. The organic layer was washed with sat. brine, dried and concentrated under vacuum to give a darkbrown oil. The oil was dissolved in methanol (400 ml) and conc. hydrochloric acid (80 ml) was added. The solution was stirred overnight at room temperature, neutralised with 5 M

sodium hydroxide and extracted three times with ether. The combined extracts were filtered through a column with  $\text{SiO}_2$  and the eluate concentrated under vacuum to give the methylated anthranilic ester (20 g). The ester was dissolved in dichloromethane (400 ml) and cooled on an ice-bath. Ethyl malonyl chloride (21 g, 0.14 mol) was added and then, after 30 minutes, triethylamine (22 ml, 0.16 mol). After being stirred for 1 hour at room temperature the cloudy mixture was washed with 0.5 M hydrochloric acid and then bicarbonate. The organic phase was carefully dried and concentrated under vacuum. The residue was then dissolved in dry ethanol (200 ml) and sodium methoxide (17 g, 0.32 mol) was added. The mixture was stirred for 1 hour and water was added (300 ml). The solution was washed with ethyl acetate and then the aqueous solution was acidified with conc. hydrochloric acid. The precipitate was collected by filtration and dried under vacuum to give the title compound as grey crystals (17 g, overall yield 41 %).

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.45 (3H, t), 3.58 (3H, s), 4.48 (2H, q), 6.17 (2H, s), 6.71 (1H, d), 7.14 (1H, d).

#### *Example 5.*

##### 5-Ethyl isatoic anhydride.

A mixture of chloral hydrate (59.3 g, 0.36 mol), water (700 ml), and sodium sulphate (85.8 g, 0.60 mol), was heated to 50°C. When 50°C was reached, sequentially a mixture of 3-ethyl-aniline (40.8 g, 0.33 mol), water (700 ml) and conc. hydrochloric acid (33.6 ml) and a mixture of hydroxylamine hydrochloride (74.8 g, 1.04 mol) and water (330 ml) were added. The resulting mixture was heated to 80°C during 30 minutes and kept for another 10 minutes at this temperature before the reaction mixture was cooled on an ice-bath. The resulting precipitate was filtered off, washed with water and dried in vacuum over  $\text{P}_2\text{O}_5$  to give an isonitrosoacetanilide (36.6 g), yield 58 %. The isonitrosoacetanilide (10.0 g, 0.05 mol), was added portionwise to a mixture of water (9 ml) and conc. sulphuric acid (60 ml) prewarmed to 50°C, maintaining the temperature between 50-55 °C. When the addition was completed, the mixture was heated to 80°C and kept at this temperature for 10 minutes. The reaction mixture was then cooled on an ice-bath and poured on 10-12 times the reaction volume of crushed ice. The mixture was then left standing for about one hour. The water suspension was extracted with dichloromethane which was dried and evaporated resulting in an mixture of the two

analogues 4-ethyl and 6-ethyl isatins approximately 0.68:1 (7.6 g), yield 84%.

The mixture of the two isomers was dissolved in aqueous sodium hydroxide and the solution was filtered through celite and then acidified to pH 4. The 4-analogue was at this pH extracted into dichloromethane which was dried and evaporated to give the pure 4-ethyl isatin (3.1 g), yield 34%.

4-Ethyl isatin (3.1 g, 0.018 mol) was added to a mixture of conc. sulphuric acid (45 µl) in acetic acid (14 ml). The suspension was warmed to 30°C, hydrogen peroxide 35% (2.2 ml) was added and after the addition the temperature was raised to 65°C. After being heated for 3 hours, the mixture was cooled and the precipitate filtered off, washed with water and dried to give the title compound (1.7 g), yield 48%.

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 1.12 (3H, t), 3.02 (2H, q), 6.98 (1H, d), 7.05 (1H, d), 7.58 (1H, t), 11.6 (1H, broad).

*Example 6.*

1,2-Dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxylic acid.

While cooling, 10 ml of conc. hydrochloric acid was added to 30 ml of acetic anhydride. To this solution, 1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxylic acid ethyl ester (10.5 g, 38 mmol) was added and the mixture heated at 80°C for 14 hours. The mixture was cooled to room temperature and the crystalline product was filtered off, washed with cold methanol and dried to yield the title compound (7.2 g), yield 77 %.

<sup>1</sup>H NMR ((CDCl<sub>3</sub>)) δ 3.73 (3H, s) 4.02 (3H, s), 6.82 (1H, d), 7.02 (1H, d), 7.62 (1H, t).

*Example 7.*

N-Methyl-N-phenyl-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide (not included in the claims) (Method A).

N-Methylaniline (2.7g, 0.025 mol) was dissolved in 80 ml of toluene and about 30 ml of the solvent was distilled off in order to obtain a dry solution. 1,2-Dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxylic acid ethyl ester (2.7 g, 0.01 mol) was added to the boiling solution. The ethanol formed during the reaction was distilled off together with some

toluene for about 4 hours. The mixture was cooled to room temperature. The precipitate was collected, washed with cold toluene and hexane and dried to give the title compound (2.8 g), yield 83 %.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.49 (3H, s), 3.50 (3H, s), 4.03 (3H, s), 6.66 (1H, d), 6.86 (1H, d), 7.08-7.48 (6H, m).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  29.7 (CH<sub>3</sub>), 36.8 (CH<sub>3</sub>), 56.8 (CH<sub>3</sub>), 103.3 (CH), 104.2 (C), 108.4 (CH), 110.2 (C), 126.2 (CH), 127.2 (CH), 128.6 (CH), 131.4 (CH), 141.2 (C), 143.6 (C), 157.0 (C), 157.4 (C), 160.3 (C), 165.1 (C).

ESI MS/MS  $[\text{M}+\text{H}]^+$  339, fragment 232.

In essentially the same manner the following compounds were obtained from the corresponding starting materials:

N-methyl-N-phenyl-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide, (not included in the claims)

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.38 (3H, s), 3.52 (3H, s), 7.08-7.34 (7H, m), 7.43 (1H, t).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  29.9 (CH<sub>3</sub>), 38.5 (CH<sub>3</sub>), 104.7 (C), 112.8 (C), 113.3 (CH), 125.5 (CH), 125.6 (CH), 126.8 (CH), 128.7 (CH), 131.8 (CH), 132.9 (C), 142.6 (C), 143.9 (C), 158.0 (C), 166.1 (C), 169.3 (C).

ESI MS/MS  $[\text{M}+\text{H}]^+$  343, fragments 236 and 108.

N-ethyl-N-(3-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-ethyl-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(4-chloro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.38 (3H, s), 3.45 (3H, s), 7.12-7.28 (6H, m), 7.45 (1H, t).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  30.0 (CH<sub>3</sub>), 38.4 (CH<sub>3</sub>), 104.5 (C), 112.6 (C), 113.4 (CH), 125.6 (CH), 127.0 (CH), 128.9 (CH), 131.9 (CH), 132.4 (C), 132.8 (CH), 142.5 (C), 142.6 (C), 158.0 (C), 166.0 (C), 169.2 (C).

ESI MS/MS  $[\text{M}+\text{H}]^+$  377, fragments 236 and 142.



N-ethyl-N-(4-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.18 (3H, t), 3.33 (3H, s), 3.74 (3H, s), 3.90 (2H, q, broad), 6.73 (2H, d), 7.05-7.15 (3H, m), 7.22 (1H, d), 7.39 (1H, t).

$^{13}\text{C NMR}$  ( $\text{CDCl}_3$ )  $\delta$  12.4 (CH<sub>3</sub>), 31.1 (CH<sub>3</sub>), 45.6 (CH<sub>2</sub>), 55.4 (CH<sub>3</sub>), 109.5 (C), 111.5 (C), 114.2 (CH), 115.2 (CH), 126.2 (CH), 127.9 (CH), 130.4 (C), 132.2 (CH), 133.1 (C), 141.7 (C), 159.2 (C), 159.3 (C), 160.1 (C), 166.7 (C).

ESI MS/MS  $[\text{M}+\text{H}]^+$  387, fragments 236 and 152.

N-methyl-N-(4-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  3.37 (3H, broad signal), 3.43 (3H, s), 3.75 (3H, s), 6.75 (2H, broad signal), 7.14 (3H, broad signal), 7.22 (1H, d), 7.40 (1H, t).

$^{13}\text{C NMR}$  ( $\text{CDCl}_3$ )  $\delta$  30.0 (CH<sub>3</sub>), 38.5 (CH<sub>3</sub>), 55.4 (CH<sub>3</sub>), 105.4 (C), 112.8 (C), 113.4 (CH), 113.9 (CH), 125.5 (CH), 127.0 (CH), 131.7 (CH), 132.7 (C), 136.8 (C), 142.6 (C), 158.1 (C), 158.3 (C), 164.9 (C), 169.1 (C).

ESI MS/MS  $[\text{M}+\text{H}]^+$  373, fragments 236 and 138.

N-ethyl-N-(3-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.20 (3H, t), 3.33 (3H, s), 3.95 (2H, q), 6.84-6.98 (3H, m), 7.11-7.20 (2H, m), 7.23 (1H, d), 7.42 (1H, t).

$^{13}\text{C NMR}$  ( $\text{CDCl}_3$ )  $\delta$  12.9 (CH<sub>3</sub>), 29.9 (CH<sub>3</sub>), 45.8 (CH<sub>2</sub>), 104.7 (C), 112.7 (C), 113.4 (CH), 113.8+114.0 (CH), 113.9+114.1 (CH), 122.3+122.4 (CH), 125.6 (CH), 129.5+129.6 (CH), 131.9 (CH), 132.8 (CH), 142.7 (C), 143.7+143.8 (C), 158.0 (C), 161.4+163.4 (C), 165.9 (C), 168.8 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS  $[\text{M}+\text{H}]^+$  375, fragments 236 and 140.

N-methyl-N-(2-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.47 (3H, s), 3.53 (3H, s), 4.03 (3H, s), 6.68 (1H, d), 6.88-6.96 (2H, m), 7.02-7.07 (1H, m), 7.12-7.17 (1H, m), 7.42-7.49 (2H, m).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 30.7 (CH<sub>3</sub>), 36.8 (CH<sub>3</sub>), 57.1 (CH<sub>3</sub>), 104.3 (C), 104.4 (CH), 107.2 (CH), 109.2 (C), 116.4+116.6 (CH), 124.3+124.3 (CH), 128.7 (CH), 129.9+130.0 (C), 129.9+130.0 (CH), 132.9 (CH), 141.1 (C), 157.4 (C), 157.4 (C), 156.8+158.8 (C), 160.3 (C), 167.0 (C).

ESI MS/MS [M+H]<sup>+</sup> 357, fragment 232.

N-methyl-N-(3-chloro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.40 (3H, s), 3.50 (3H, s), 4.02 (3H, s), 6.67 (1H, d, broad), 6.90 (1H, d, broad), 7.1 (2H, broad), 7.28 (1H, broad), 7.38 (1H, broad), 7.43 (1H, t, broad).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 29.8 (CH<sub>3</sub>), 36.8 (CH<sub>3</sub>), 57.0 (CH<sub>3</sub>), 103.5 (CH), 104.3 (C), 108.6 (CH), 109.9 (C), 124.7 (CH), 126.5 (CH), 127.5 (CH), 129.7 (CH), 131.7 (CH), 133.9 (C), 141.4 (C), 144.8 (C), 157.2 (C), 157.7 (C), 160.3 (C), 165.0 (C).

ESI MS/MS [M+H]<sup>+</sup> 373, fragment 232.

N-ethyl-N-(3-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.22 (3H, t), 3.50 (3H, s), 3.92 (2H, broad signal), 4.02 (3H, s), 6.66 (1H, d), 6.81-6.92 (2H, m), 7.08-7.19 (3H, m), 7.41 (1H, t).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 13.1 (CH<sub>3</sub>), 29.8 (CH<sub>3</sub>), 43.9 (CH<sub>2</sub>), 56.9 (CH<sub>3</sub>), 103.4 (CH), 104.3 (C), 108.6 (CH), 110.4 (C), 114.5+114.7 (CH), 123.4 (CH), 129.6+129.7 (CH), 131.6 (CH), 141.4 (C), 143.5 (C), 157.2 (C), 157.4 (C), 160.3 (C), 161.4+163.3 (C), 164.4 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS [M+H]<sup>+</sup> 371, fragment 232.

N-methyl-N-(4-chloro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.46 (3H, s), 3.52 (3H, s), 4.05 (3H, s), 6.69 (1H, d), 6.92 (1H, d), 7.10-7.38 (4H, dd), 7.45 (1H, t).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  29.8 (CH<sub>3</sub>), 36.8 (CH<sub>3</sub>), 56.8 (CH<sub>3</sub>), 103.4 (CH), 104.2 (C), 108.6 (CH), 110.0 (C), 127.6 (CH), 128.9 (CH), 131.6 (CH), 132.8 (C), 141.3 (C), 142.2 (C), 157.1 (C), 157.5 (C), 160.3 (C), 165.0 (C).

ESI MS/MS  $[\text{M}+\text{H}]^+$  373, fragment 232.

N-ethyl-N-(2-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-trifluoromethyl-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(4-chloro-phenyl)-1,2-dihydro-4-hydroxy-5-trifluoromethyl-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.40 (3H, s), 3.48 (3H, s), 7.08-7.25 (4H, m), 7.48 (1H, d), 7.65 (1H, t), 7.69 (1H, t).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  30.1 (CH<sub>3</sub>), 38.7 (CH<sub>3</sub>), 103.8 (C), 112.7 (C), 113.4 (C), 118.7 (CH), 121.9+121.9+122.0+122.0 (CH), 120.3+122.4+124.6+126.8 (C), 127.0 (CH), 127.8+128.0+128.3+128.5 (C), 128.9 (CH), 131.6 (CH), 132.4 (C), 142.3 (C), 142.6 (C), 157.7 (C), 166.3 (C), 169.9 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS  $[\text{M}+\text{H}]^+$  411, fragments 270 and 142.

N-methyl-N-(4-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-trifluoromethyl-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(4-chloro-phenyl)-1,2-dihydro-4-hydroxy-5-trifluoromethoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(4-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-trifluoromethoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-phenyl-1,2-dihydro-4-hydroxy-5-chloro-6-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide, (not included in the claims)

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.38 (3H, s, broad), 3.52 (3H, s), 3.96 (3H, s), 7.14-7.23 (2H, m), 7.23-7.30 (5H, m).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  29.7 (CH<sub>3</sub>), 38.3 (CH<sub>3</sub>), 57.2 (CH<sub>3</sub>), 113.6 (CH), 113.7 (C), 116.8 (CH), 120.3 (C), 125.8 (CH), 126.9 (CH), 128.7 (CH), 136.5 (C), 143.9 (C), 150.9 (C), 158.0 (C), 165 (C), 168.9 (C).

ESI MS/MS  $[\text{M}+\text{H}]^+$  373, fragments 266 and 108.

N-methyl-N-(4-chloro-phenyl)-1,2-dihydro-4-hydroxy-5,6-methylenedioxy-1-methyl-2-oxo-quinoline-3-carboxamide.

*Example 8.*

N-Methyl-N-(4-trifluoromethyl-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide (Method B).

To an ice-cold solution of 1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxylic acid (8 g, 0.032 mol), triethylamine (15.5 ml, 0.11 mol) and 4-trifluoromethyl-N-methylaniline (6.1 g, 0.035 mol) in 150 ml of methylene chloride was added dropwise during 0.5 hours a solution of thionyl chloride (3.0 ml, 0.042 mol) in 10 ml of methylene chloride. The stirring was continued at 4°C for 4 hours. The solution was diluted with 10 ml of methylene chloride, washed with cold 1 M sulphuric acid and then extracted with 1 M sodium hydroxide. The pH of the aqueous phase was adjusted to 8-8.5, clarified by filtration and then acidified with hydrochloric acid to pH 4. On standing a crystalline precipitate was formed which was filtered off, washed with water and dried to give the title compound (8.5 g) yield 65 %.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.48 (3H, s), 3.54 (3H, s), 4.06 (3H, s), 6.70 (1H, d), 6.94 (1H, d), 7.46 (1H, t), 7.50 (4H, broad signal).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  29.8 (CH<sub>3</sub>), 36.9 (CH<sub>3</sub>), 56.9 (CH<sub>3</sub>), 103.5 (CH), 104.2 (C), 108.7 (CH), 109.5 (C), 117.3+121.7+126.0+130.3 (C), 125.8+125.9+125.9+126.0 (CH), 126.3

(CH), 127.9+128.4+128.9+129.4 (C), 131.8 (CH), 141.4 (C), 146.7 (C), 157.2 (C), 158.0 (C), 160.3 (C), 165.0 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS  $[M+H]^+$  407, fragment 232.

In essentially the same manner the following compounds were obtained from the corresponding starting materials:

N-ethyl-N-(4-trifluoromethyl-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.22 (3H, t), 3.28 (3H, s), 3.99 (2H, q), 7.13 (1H, d), 7.23-7.32 (3H, m), 7.40-7.51 (3H, m).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  13.0 (CH<sub>3</sub>), 29.8 (CH<sub>3</sub>), 45.8 (CH<sub>2</sub>), 104.0 (C), 112.7 (C), 113.5 (CH), 120.6+122.8+124.9+127.1 (C), 125.7 (CH), 125.7+125.7+125.8+125.8 (CH), 126.7 (CH), 128.3+128.6+128.8+129.1 (C), 132.1 (CH), 133.0 (C), 142.8 (C), 145.6 (C), 157.9 (C), 166.8 (C), 169.1 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS  $[M+H]^+$  425, fragments 236 and 190.

N-ethyl-N-(4-trifluoromethyl-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.22 (3H, t), 3.51 (3H, s), 3.93 (2H, q), 4.02 (3H, s), 6.67 (1H, d), 6.91 (1H, d), 7.43 (1H, t), 7.46-7.52 (4H, m).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  13.2 (CH<sub>3</sub>), 29.8 (CH<sub>3</sub>), 44.1 (CH<sub>2</sub>), 56.9 (CH<sub>3</sub>), 103.5 (CH), 104.3 (C), 108.7 (CH), 110.0 (C), 120.7+122.9+125.0+127.2 (C), 125.9+125.9 (CH), 127.7 (CH), 128.9+129.2+129.4+129.7 (C), 131.8 (CH), 141.5 (C), 145.3 (C), 157.2 (C), 157.8 (C), 160.3 (C), 164.4 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS  $[M+H]^+$  421, fragments 232 and 206.

N-methyl-N-(4-trifluoromethoxy-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.33 (3H, s), 3.45 (3H, s), 6.62 (1H, broad), 6.83 (1H, broad), 6.98-7.17 (2H, m, broad), 7.20 (1H, d), 7.37 (1H, t, broad).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 29.9 (CH<sub>3</sub>), 37.3 (CH<sub>3</sub>), 103.3 (C), 104.7+104.9+105.1 (CH), 110.5+110.7 (CH), 112.7 (C), 113.3 (CH), 125.7 (CH), 128.1 (C), 128.6 (CH), 132.1 (CH), 133.3 (C), 142.8 (C), 157.8 (C), 156.9+157.0+158.9+159.0 (C), 160.6+160.6 (C), 167.4 (C), 170.4 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS [M+H]<sup>+</sup> 379, fragments 236 and 144.

N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide,

N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.40 (3H, s), 3.51 (3H, s), 4.02 (3H, s), 6.60-6.63 (1H, m), 6.63 (1H, d), 6.73-6.79 (1H, m), 6.90 (1H, d), 7.38-7.46 (2H, m).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 29.9 (CH<sub>3</sub>), 36.0 (CH<sub>3</sub>), 56.9 (CH<sub>3</sub>), 103.5 (CH), 104.2 (C), 104.4+104.6+104.6+104.8 (CH), 108.6 (CH), 109.2 (C), 110.8+ 110.9+111.0+111.0 (CH), 127.3+127.3+127.4+127.4 (C), 130.0+130.1 (CH), 131.8 (CH), 141.4 (C), 157.2 (C), 157.3+157.4+159.3+159.4 (C), 158.5 (C), 160.3 (C), 160.7+160.8+162.6+162.7 (C), 165.5 (C); some peaks are multiplets due to F-coupling.

ESI MS/MS [M+H]<sup>+</sup> 375, fragment 232.

N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide,

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.48 (3H, s), 3.64 (3H, s), 4.10 (3H, s), 6.60-7.30 (5H, m), 7.63 (1H, t).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 31.0 (CH<sub>3</sub>), 37.2 (CH<sub>3</sub>), 57.2 (CH<sub>3</sub>), 104.4 (C), 105.0 (CH), 105.7 (C), 109.5 (CH), 115.2+115.6 (CH), 116.8+116.9 (CH), 117.2+117.3+117.5+117.7 (CH), 129.8+130.0+130.0+130.2 (C), 133.9 (CH), 141.0 (C), 151.9+155.8 (C), 157.6 (C),

155.8+159.6 (C), 161.4 (C), 161.7 (C), 167.6 (C); major form; some peaks are multiplets due to F-coupling.

ESI MS/MS  $[M+H]^+$  375, fragment 232.

N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5,6-methylenedioxy-1-methyl-2-oxo-quinoline-3-carboxamide.

### *Pharmacological methods*

#### Acute experimental autoimmune encephalomyelitis (aEAE).

SJL/N female mice, 8 weeks of age, were used for the experiments. Mouse spinal cord homogenate (MSCH) was obtained from 8 to 12 weeks-old C57Bl/6 female mice. The tissue was homogenised on ice and diluted in cold PBS. Incomplete Freund's containing 1 mg/ml *M. tuberculosis hominis* H37Ra was emulsified with an equal volume of MSCH to give a final concentration of 10 mg/ml of MSCH. The inoculum volume of 0.1 ml was injected intradermally at the base of the tail. Pertussis toxin was injected i.p. at day 0 and 3 after immunization. Treatment was given per os daily either at day 3 to 12 post-immunization or days 3 to 7 and 10 to 12. Control animals received saline. The animals, eight per dose group, were scored for clinical signs of paralytic disease on a scale from 0 to 5 in the following way: 0, normal; 1, limp tail; 2, hind limb paresis; 3 hind limb paralysis and limp foreleg; 4, bilateral hind and fore limb paralysis; 5, death. Clinical scores were monitored at day 7 and daily from day 9 until the end of the experiment at day 14. Treatment effects were calculated as percent inhibition of clinical scores compared to saline treated controls.

#### Collagen induced arthritis.

DBA/1 male mice between 8 to 10 weeks of age were used for the experiments. On day 0 the mice were immunized intradermally at the base of the tail with bovine type II collagen (100 µg/mouse) in Freund's complete adjuvant. The treatment was given per os daily on days 3 to 7, 10 to 14, 17 to 21, 24 to 28 and 31 to 35. Fifteen days after immunization mice were inspected for signs of arthritis. The animals were inspected three times a week. Every second or third day individual paws of the arthritic animals were scored by a scale from 0-4 (0 = no arthritis, 1 = arthritis in one of the interphalangeal, metatarsophalangeal or intercarpal joints.

2 = two arthritic joints, 3 = three arthritic joints, 4 = as in 3 but with more severe redness and swelling of the paw). The score for each paw was added to give a maximal attainable score of 16 for each mouse.

#### Ovalbumin-induced lung inflammation.

C57Bl/6 female mice between 10 to 14 weeks of age were used for the experiments, 10 mice /group. The mice were sensitized with ovalbumin (OA) in aluminium hydroxide in a volume of 0.2 ml inoculated ip. Treatment was given at day 0 to day 16. Control mice received saline. Fourteen days after the OA sensitization mice were exposed for 20 minutes to an aerosol of 1.5% w/v of OA in saline produced by a nebulizer. Vehicle-challenged control mice were exposed to saline. Seventy-two hours after OA/vehicle challenge, mice were anaesthetised and bronchoalveolar lavage was performed by instilling 0.5 ml ice-cold phosphate buffered saline (PBS) into the lungs twice. Total cell counts were determined and differential counts were made based on identification of eosinophils, monocytes/alveolar macrophages, lymphocytes and neutrophils. Eosinophil infiltration into the lung tissue was evaluated by histochemical methods on frozen lung sections using diaminobenzidine tetrahydrochloride (DAB).

#### *Teratogenic effects in the rat.*

The compounds were administrated subcutaneously to female rats during pregnancy, i.e., day 8 to 14 of pregnancy. The rats were caesarean sectioned and necropsied on day 20 after fertilisation. The foetuses were examined for external and internal abnormalities.

#### *Beagle Pain Syndrome (BPS).*

The compounds were administrated intravenously to beagle dogs. The dosage was given for five consecutive days. The dogs were evaluated for clinical and laboratory signs of the pain syndrome, e.g., fever, increased erythrocyte sedimentation rate (ESR), alkaline phosphate (AP), induction of acute phase proteins and vasculitis

Among preferred compounds are N-methyl-N-(4-trifluoromethyl-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide and N-methyl-N-(2,4-difluoro-



phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide hereinafter called Compound A and B, respectively. N-Methyl-N-phenyl-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide and roquinimex are included as reference compounds hereinafter called Compound C and D, respectively:

aEAE inhibition

Dose, mg/kg p.o.	% aEAE Inhibition			
	Compound A (invention)	Compound B (invention)	Compound C	Compound D
0.2	66	59	92	35
1	86	96	100	40
5	99	100		69

Teratogenicity in the rat

Dose, mg/kg p.o.	% malformed fetuses			
	Compound A (invention)	Compound B (invention)	Compound C	Compound D
6	4	0	37	not tested
10	not tested	not tested	not tested	9
30	2			30

Effective quantities of the compounds of formula (I) are preferably administered to a patient in need of such treatment according to usual routes of administration and formulated in usual pharmaceutical compositions comprising an effective amount of the active ingredient and a suitable pharmaceutically acceptable carrier. Such compositions may take a variety of forms, e.g. solutions, suspensions, emulsions, tablets, capsules, and powders prepared for oral administration, aerosols for inhalations, sterile solutions for parental administration, suppositories for rectal administration or suitable topical formulations. Conventional procedures for the selection and preparation of suitable pharmaceutical formulations are

described, for example, in "Pharmaceuticals - The Science of Dosage Form Design", M.B. Aulton, Churchill Livingstone, 1988.

A suitable daily dose for use in the treatment of MS is contemplated to vary between 0.0005 mg/kg to about 10 mg/kg body weight, in particular between 0.005 mg/kg to 1 mg/kg body weight, depending upon the specific condition to be treated, the age and weight of the specific patient, and the specific patient's response to the medication. The exact individual dosage, as well as the daily dosage, will be determined according to standard medical principles under the direction of a physician.

Various additives to enhance the stability or ease of administration of the drug are contemplated. The pharmaceutical composition may also contain additional therapeutically useful substances other than a compound of formula (I).

#### *References*


1. Talal, N.: Autoimmune diseases. In: Roitt, I.M. and Delves, P.J. (eds.) *Encyclopedia of Immunology*, pp. 195-198. Academic Press, 1992.
2. Prineas, J.W.: The neuropathology of multiple sclerosis. In: Koetsier, J.C. (ed.) *Handbook of Clinical Neurology*, pp. 213-257. Elsevier Science Publ., Amsterdam, 1985.
3. Tarkowski, A., Gunnarsson, K., Nilsson, L.-Å., Lindholm, L. and Stålhandske, T. Successful treatment of autoimmunity in MRL/l mice with LS2616, a new immunomodulator. *Arthritis Rheum.* 29(11):1405-1409, 1986.
4. Larsson, E.-L., Joki, A.-L. and Stålhandske, T. Mechanism of action of the new immunomodulator LS2616 on T-cell responses. *Int J Immunopharmacol* 9(4):425-31, 1987.
5. Wanders, A., Larsson, E., Gerdin, B. and Tufveson G. Abolition of the effect of cyclosporine on rat cardiac allograft rejection by the new immunomodulator LS-2616 (Linomide). *Transplantation* 47(2):216-217, 1989.
6. Kalland, T. Regulation of natural killer progenitors: studies with a novel immunomodulator with distinct effects at the precursor level. *J Immunol* 144(11): 4472-4476, 1990.
7. Gonzalo, J.A., González-García, A., Kalland, T., Hedlund, G., Martinez, C. and Kroemer, G. Linomide, a novel immunomodulator that prevents death in four models of septic shock.

Eur J Immunol 23:2372-2374, 1993.

8. Karussis, D.M., Lehmann, D., Slavin, S. et al. Treatment of chronic-relapsing experimental autoimmune encephalomyelitis with the syntethic immunomodulator Linomide (quinoline-3-carboxamide). Proc Natl Acad Sci USA 90: 6400-6404, 1993.
9. Gonzalo, J.A., González-Garcia, A., Kalland, T. et al. Linomide inhibits programmed cell death of peripheral T cells in vivo. Eur J Immunol. 24: 48-52, 1994.
10. Gross, D.J., Sidi, H., Weiss, L., Kalland, T., Rosenmann, E. and Slavin, S. Prevention of diabetes mellitus in non-obese diabetic mice by Linomide, a novel immunomodulating drug. Diabetologia 37: 1195-1201, 1994.
11. Karussis, D.M., Lehmann, D., Brenner, T. et al. Immunomodulation of experimental autoimmune myasthenia gravis with Linomide. J Neuroimmunol 55(2): 187-193, 1994.
12. Bai, X.F., Shi, F.D., Zhu, J., Xiao, B.G., Hedlund, G. and Link, H. Linomide-induced suppression of experimental autoimmune neuritis is associated with down-regulated macrophage functions. J Neuroimmunol 76:177-184 1997.
13. Karussis, D.M. Meiner, Z., Lehmann, D. et al. Treatment of secondary progressive multiple sclerosis with the immunomodulator Linomide. Neurology 47: 341-346, 1996.
14. Andersen, O., Lycke, J., Tolleson, P.O. et al. Linomide reduces the rate of active lesions in relapsing-remitting multiple sclerosis. Neurology 47: 895-900, 1996.
15. Kelly, D.F., Grimsell, C.S.G. and Kenyon, C.J. Polyarteritis in the dog: A case report. Vet Record 92: 363-366, 1973.
16. Harcourt, R.A. Polyarterites in a colony of beagles. Vet Record 102: 519-522, 1978.

**We claim:**

### 1. Compounds of general formula (I)



wherein

R is selected from methyl, ethyl, n-propyl, iso-propyl, n-butyl and allyl;

R' is selected from methyl, methoxy, fluoro, chloro, bromo, trifluoromethyl, and  $\text{OCH}_2\text{F}$ .

wherein  $x = 0 - 2$ ,

$y = 1 - 3$  with the proviso that

 $x + y = 3;$ 

R'' is selected from hydrogen, fluoro and chloro, with the proviso that R'' is selected from fluoro and chloro only when R' is selected from fluoro and chloro;

$R_4$  is selected from hydrogen and pharmaceutically acceptable inorganic and organic cations;

$R_5$  is selected from ethyl, n-propyl, iso-propyl, methoxy, ethoxy, chloro, bromo, trifluoromethyl,  $OCH_2F_y$ , and  $OCH_2CH_2F_y$

wherein  $x = 0 - 2$ ,

$y = 1 - 3$  with the proviso that

$$x + y = 3;$$

$R_6$  is hydrogen; or


$R_5$  and  $R_6$  taken together are methylenedioxy;

and any tautomer thereof.

2. Compounds according to claim 1 wherein the pharmaceutically acceptable inorganic cations are derived from sodium, potassium and calcium, and the organic cations are derived from monoethanolamine, diethanolamine, dimethylaminoethanol, morpholine and the like;


3. Compounds according to claim 1 and 2 wherein  $R_5$  is selected from ethyl, methoxy, chloro, and trifluoromethyl.
4. Compounds according to claim 1 and 2 wherein  $R_5$  and  $R_6$  taken together are methylenedioxy.
5. Compounds according to any of the preceding claims wherein R is selected from methyl and ethyl.
6. Compounds according to any of the preceding claims wherein  $R'$  is selected from, methoxy, fluoro, chloro, and trifluoromethyl, when  $R''$  is hydrogen.
7. Compounds according to any of the preceding claims wherein  $R''$  is selected from *meta*- and *para*-fluoro provided that  $R'$  is *ortho*-fluoro.
8. The compound according to claims 1 and 2, N-ethyl-N-(3-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide.
9. The compound according to claims 1 and 2, N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide.
10. The compound according to claims 1 and 2, N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide.
11. The compound according to claims 1 and 2, N-ethyl-N-(3-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-ethyl-1-methyl-2-oxo-quinoline-3-carboxamide.
12. The compound according to claims 1 and 2, N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide.

13. The compound according to claims 1 and 2, N-methyl-N-(4-trifluoromethyl-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide.
14. The compound according to claims 1 and 2, N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5,6-methylenedioxy-1-methyl-2-oxo-quinoline-3-carboxamide.
15. The compounds according to any of the preceding claims to be used as therapeuticum.
16. Pharmaceutical compositions containing as active ingredient a compound having the general formula (I) together with a pharmaceutically carrier.
17. Pharmaceutical compositions according to claim 16 containing other pharmacologically active substances.
18. Pharmaceutical compositions according to claims 16 and 17 to be used as therapeuticum in a daily dose of the active substance of 0.0005 mg/kg to about 10 mg/kg body weight, in particular 0.005 to 1 mg/kg body weight.
19. A process for the manufacturing of a compound of the general formula (I) wherein R, R', R'', R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are as defined above, by
- (A) reacting an ester derivative of quinoline carboxylic acid of formula (II)




with an aniline of formula (III), in a suitable solvent such as toluene, xylene, or the like, or

(B) reacting a quinoline carboxylic acid of the general formula (IV) with an aniline of the general formula (III),



using a suitable coupling reagent, preferably a carbodiimide or thionyl chloride in the presence of triethylamine and a suitable solvent such as dichloromethane.

20. A method of treating a mammal suffering from pathologic inflammation and autoimmunity comprising administering to said mammal a compound having the general formula (I)



wherein

R is selected from methyl, ethyl, n-propyl, iso-propyl, n-butyl and allyl;

R' is selected from methyl, methoxy, fluoro, chloro, bromo, trifluoromethyl, and  $\text{OCH}_x\text{F}_y$ ,

wherein  $x = 0 - 2$ ,

$y = 1 - 3$  with the proviso that

$x + y = 3$ ;

R'' is selected from hydrogen, fluoro and chloro, with the proviso that R'' is selected from fluoro and chloro only when R' is selected from fluoro and chloro;

R<sub>4</sub> is selected from hydrogen and pharmaceutically acceptable inorganic and organic cations;

R<sub>5</sub> is selected from ethyl, n-propyl, iso-propyl, methoxy, ethoxy, chloro, bromo,

trifluoromethyl,  $\text{OCH}_x\text{F}_y$ , and  $\text{OCH}_2\text{CH}_x\text{F}_y$

wherein  $x = 0 - 2$ ,  
 $y = 1 - 3$  with the proviso that  
 $x + y = 3$ ;

$R_4$  is hydrogen; or  
 $R_5$  and  $R_6$  taken together are methylenedioxy;  
and any tautomer thereof.

21. The method according to claim 20 wherein the pharmaceutically acceptable inorganic cations are derived from sodium, potassium and calcium, and the organic cations are derived from monoethanolamine, diethanolamine, dimethylaminoethanol, morpholine and the like;

22. The method according to claim 20 and 21 wherein  $R_5$  is selected from ethyl, methoxy, chloro, and trifluoromethyl.

23. The method according to claim 20 and 21 wherein  $R_5$  and  $R_6$  taken together are methylenedioxy.

24. The method according to claim 20 and 21 wherein  $R$  is selected from methyl and ethyl.

25. The method according to claim 20 and 21 wherein  $R'$  is selected from, methoxy, fluoro, chloro, and trifluoromethyl when  $R''$  is hydrogen.

26. The method according to claim 20 and 21 wherein  $R''$  is selected from *meta'*- and *para*-fluoro provided that  $R'$  is *ortho*-fluoro.

27. The method according to claim 20 and 21, N-ethyl-N-(3-fluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide.

28. The method according to claim 20 and 21, N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide.



29. The method according to claim 20 and 21, N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide.
30. The method according to claim 20 and 21, N-ethyl-N-(3-methoxy-phenyl)-1,2-dihydro-4-hydroxy-5-ethyl-1-methyl-2-oxo-quinoline-3-carboxamide.
31. The method according to claim 20 and 21, N-methyl-N-(2,5-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide.
32. The method according to claim 20 and 21, N-methyl-N-(4-trifluoromethyl-phenyl)-1,2-dihydro-4-hydroxy-5-methoxy-1-methyl-2-oxo-quinoline-3-carboxamide.
33. The method according to claim 20 and 21, N-methyl-N-(2,4-difluoro-phenyl)-1,2-dihydro-4-hydroxy-5,6-methylenedioxy-1-methyl-2-oxo-quinoline-3-carboxamide.
34. The method according to any of claims 20 to 33 of treating mammals suffering from multiple sclerosis (MS).
35. The method according to any of claims 20 to 33 of treating mammals suffering from insulin-dependent diabetes mellitus (IDDM).
36. The method according to any of claims 20 to 33 of treating mammals suffering from systemic lupus erythematosus (SLE).
37. The method according to any of claims 20 to 33 of treating mammals suffering from rheumatoid arthritis (RA).
38. The method according to any of claims 20 to 33 of treating mammals suffering from inflammatory bowel disease (IBD).

39. The method according to any of claims 20 to 33 of treating mammals suffering from psoriasis.
40. The method according to any of claims 20 to 33 of treating mammals suffering from inflammatory respiratory disorder, such as asthma.
41. The method according to any of claims 20 to 33 of treating mammals suffering from atherosclerosis.
42. The method according to any of claims 20 to 33 of treating mammals suffering from stroke.
43. The method according to any of claims 20 to 33 of treating mammals suffering from Alzheimer's disease.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE 99/01270

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 215/56, C07D 215/22, C07D 215/18, A61K 31/47  
According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D, A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0059698 A1 (AB LEO), 8 Sept 1982 (08.09.82), page 3, line 10 - line 15 --	1-43
X	GB 2290786 A (FUJISAWA PHARMACEUTICAL CO., LTD.), 10 January 1996 (10.01.96), page 10 - page 11 --	1-43
X	WO 9524395 A1 (FUJISAWA PHARMACEUTICAL CO., LTD.), 14 Sept 1995 (14.09.95), page 25 - page 26 --	1-43
X	WO 9218483 A1 (FUJISAWA PHARMACEUTICAL CO., LTD.), 29 October 1992 (29.10.92), page 53 - page 54 --	1-43



Further documents are listed in the continuation of Box C.



See patent family annex.

- \* Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

21 October 1999

Date of mailing of the international search report

30-10-1999

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Nebil Gecer/ELY  
Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 99/01270

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Patent Abstracts of Japan, abstract of JP 7-224040 A (FUJISAWA PHARMACEUT CO LTD.), 22 August 1995 (22.08.95)	1-43
	--	
A	WO 9524196 A1 (PHARMACIA AB), 14 Sept 1995 (14.09.95)	1-43
	--	
A	WO 9524195 A1 (PHARMACIA AB), 14 Sept 1995 (14.09.95)	1-43
	--	
A	WO 9319756 A1 (KABI PHARMACIA AB), 14 October 1993 (14.10.93)	1-43
	--	
A	WO 9306829 A1 (KABI PHARMACIA AB), 15 April 1993 (15.04.93)	1-43
	-- -----	

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE99/01270

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 20-43  
because they relate to subject matter not required to be searched by this Authority, namely:  
See extra sheet.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE99/01270

Claims 20-43 are directed to methods of treatment of the human or animal body by therapy methods practised on the human or animal body (see PCT, Rule 39.1(iv)). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

28/09/99

International application No.

PCT/SE 99/01270

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0059698 A1	08/09/82	SE 0059698 T3 AT 15798 T AU 547849 B AU 8107682 A CA 1182455 A DK 81282 A DK 157859 B,C FI 79532 B,C FI 820728 A GR 76073 A HK 38689 A IE 52670 B JP 1721152 C JP 4009786 B JP 57171975 A NZ 199888 A PH 18520 A PT 74512 A,B US 4547511 A US 4738971 A ZA 8201210 A	15/10/85 07/11/85 09/09/82 12/02/85 04/09/82 26/02/90 29/09/89 04/09/82 03/08/84 19/05/89 20/01/88 24/12/92 21/02/92 22/10/82 31/07/85 02/08/85 01/04/82 15/10/85 19/04/88 26/01/83
GB 2290786 A	10/01/96	GB 9413150 D	00/00/00
WO 9524395 A1	14/09/95	AU 1823595 A CA 2214521 A GB 9404378 D JP 9509957 T	25/09/95 14/09/95 00/00/00 07/10/97
WO 9218483 A1	29/10/92	AU 656576 B AU 1548792 A CA 2108971 A EP 0639182 A HU 67349 A HU 9302983 D JP 6506925 T MX 9201823 A	09/02/95 17/11/92 23/10/92 22/02/95 28/03/95 00/00/00 04/08/94 01/10/92
WO 9524196 A1	14/09/95	AU 679633 B AU 2088795 A CA 2185135 A DE 69406800 D,T EP 0717819 A,B EP 0749310 A FI 963531 A IL 112908 D IL 112909 D JP 9502786 T JP 9510199 T NO 963762 A SE 9400810 D US 5704865 A US 5726183 A ZA 9501966 A	03/07/97 25/09/95 14/09/95 23/04/98 26/06/96 27/12/96 09/09/96 00/00/00 00/00/00 18/03/97 14/10/97 09/09/96 00/00/00 06/01/98 10/03/98 11/12/95

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

28/09/99

International application No.  
PCT/SE 99/01270

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9524195 A1	14/09/95	AU 690661 B	30/04/98
		AU 2088695 A	25/09/95
		AU 7712294 A	03/04/95
		CA 2185136 A	14/09/95
		EP 0751772 A	08/01/97
		FI 963532 A	06/11/96
		IL 112908 D	00/00/00
		JP 9510198 T	14/10/97
		NO 963764 A	09/09/96
		SE 9400809 D	00/00/00
		US 5728713 A	17/03/98
		ZA 9501967 A	11/12/95
WO 9319756 A1	14/10/93	AU 667362 B	21/03/96
		AU 3913193 A	08/11/93
		CA 2131831 A,C	14/10/93
		CN 1081671 A	09/02/94
		EP 0634931 A	25/01/95
		HU 69710 A	28/09/95
		HU 9402854 D	00/00/00
		JP 7506568 T	20/07/95
		NO 943723 A	05/10/94
		NZ 251494 A	27/07/97
		SE 9201076 A	07/10/93
		US 5594005 A	14/01/97
		ZA 9302450 A	20/10/93
WO 9306829 A1	15/04/93	AU 653647 B	06/10/94
		AU 2770892 A	03/05/93
		CA 2120712 A	15/04/93
		EP 0607282 A	27/07/94
		JP 6511488 T	22/12/94
		NO 941273 A	08/06/94
		PT 100937 A	31/01/94
		SE 469368 B,C	28/06/93
		SE 9102909 A	10/04/93
		US 5580882 A	03/12/96



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**